


Crossword Puzzle



© 2008 Sylvia Rayfield, Inc.

Across:

2. Assess K level.
5. Induces sleep, protect from injury, no alcohol or other CNS depressants.
6. Only this anti-inflammatory has IM, IV and oral preparation.
7. Toxicity includes blue/green halo, nausea and confusion. Keep therapeutic blood level.
10. Administer from protocol for crushing chest pain. Watch respiratory rate.
12. Anti-infective. Assess for penicillin allergy prior to administration. Undesireable effect: GI distress.
14. Taper this anti-inflammatory slowly.
17. Diuretic. Monitor for muscle cramping. Instruct to increase potassium and decrease sodium intake. Evaluate I&O and muscle cramping.
18. Anti-diabetic that must be taken with meals and have adequate hydration.
19. This once a day anti-inflammatory is used to treat arthritis.
22. Beta blocker, contraindicated with severe bradycardia.
25. Helps breathing. Evaluate lung sounds, nervousness and tremors.
26. Hold if allergic to penicillin.
27. Remember to take with food for this short acting anti-inflammatory.

Down:

1. Caution if drug abuse history. Should not be crushed or chewed, no alcohol.
3. Do not combine this alternative with prescription anti-depressants.
4. Counsel patient regarding bleeding precautions with this alternative.
8. Alternative. Avoid use with anticoagulants, improves stamina.
9. Evaluate INR when administering this injectable.
11. Memory gets better with this alternative.
13. Inhibits the enzyme necessary for cholesterol biosynthesis in the intestine and liver. It interacts with Warfarin, and requires 2-3 liters of fluid daily.
15. Anti-infective, instruct to avoid taking with food and antacid. Increase fluids.
16. Evaluate decreased headache pain with this alternative.
20. Side effects of Breast Cancer drug include hot flashes and vaginal dryness. Take once a month if you can sit up for 30 minutes. Evaluate effectiveness with bone scan.
21. Analgesic drug, causes decrease peristalsis and CNS changes.
23. Anti-hypertensive agent. Contraindicated for patients with asthma and can mask hypoglycemia.
24. CA Channel blocker. Monitor for peripheral edema.